

# New provincial and state records for Heteroptera (Hemiptera) in Canada and the United States

G.G.E. SCUDDER<sup>1</sup>

## ABSTRACT

New provincial records are provided for 52 species of Heteroptera in Canada. *Dichaetocoris piceicola* (Knight) is also reported from Alaska, and *Pagasa nigripes* Harris from Washington State.

## INTRODUCTION

Since the publication of the checklist of the Hemiptera of Canada and Alaska (Maw *et al.* 2000), new Heteroptera from Canada, and new provincial records have been published by Barnes *et al.* (2000), Henry *et al.* (2008), Jansson (2002), Kenner and Needham (2004), Paiero *et al.* (2003), Roch (2007), Schuh (2000a, 2000b, 2001, 2004a, 2004b), Schuh and Schwartz (2004, 2005), Schwartz and Scudder (2001, 2003), Schwartz and Stonedahl (2004), Scudder (2000, 2004, 2007), Scudder and Footitt (2006), Scudder and Schwartz (2001), Wheeler and Hoebeke (2004), and Wheeler *et al.* (2006). Wright (1989) and Kerzhner (1993) also published records not included in Maw *et al.* (2000).

Recent research has revealed additional new provincial records for 52 species. *Dichaetocoris piceicola* (Knight) has been found in Alaska, and *Pagasa nigripes* Harris in Washington State.

These are reported below, with Museum abbreviations as follows:

AAFCL: Agriculture and Agri-Food Canada, Lethbridge, AB (J.R. Byers).

CNC: Canadian National Collection of Insects, Agriculture and Agri-Food Canada,

Ottawa, ON (R.G. Footitt).

LM: Lyman Entomological Museum, Macdonald College, McGill University, Ste.-Anne-de-Bellevue, QC (T. Wheeler).

MU: Memorial University. St. John's, NF (D. Larson).

NSM: Nova Scotia Museum of Natural History, Halifax, NS (A. Hebda and C. Majka).

RBCM: Royal British Columbia Museum, Victoria, BC (R.A. Cannings).

SM: Saskatchewan Provincial Museum, Saskatoon, SK (R. Hooper).

UBC: Spencer Entomological Museum, Department of Zoology, University of British Columbia, Vancouver, BC (K.M. Needham).

UG: Department of Environmental Biology, University of Guelph, Guelph, ON (S.A. Marshall).

UM: J.B. Wallis Collection, University of Manitoba, Winnipeg, MB (R.E. Roughley).

UPEI: Department of Biology, University of Prince Edward Island, Charlottetown, PE (K.A. Campbell and D. Giberon).

## NEW PROVINCIAL RECORDS

The systematic order of families and higher taxa listed below, follows Maw *et al.* (2000).

### Infraorder NEPOMORPHA

#### Family CORIXIDAE

<sup>1</sup>Department of Zoology, University of British Columbia, 6270 University Boulevard, Vancouver, BC V6T 1Z4. (604) 822-3682. scudder@zoology.ubc.ca

*Arctocoris chanceae* Hungerford

A subarctic species known from Alaska and across northern Canada from Yukon to Newfoundland and Labrador (Scudder 1997; Maw *et al.* 2000). The species can be keyed using Hungerford (1948) and Brooks and Kelton (1967).

New record. **BC**: 1♂, Atlin, 10 mi S, 18.viii.1980 (R.J. Cannings) [UBC].

*Sigara compressoidea* (Hungerford)

An eastern Nearctic species, in Canada reported from Ontario east to Newfoundland (Maw *et al.* 2000), but not previously reported from Prince Edward Island. *Sigara compressoidea* is keyed in Hungerford (1948) and Tinerella and Gundersen (2005), and a dorsal colour photograph is given in the latter reference, showing the typically effaced membrane patterning.

New record. **PE**: 1♂ 1♀, Deroche Natural Protected Area, 46.42114°N 62.94082°W, kicknet, small wetland, 27.ix.2007 (K. Alexander Campbell) [UPEI].

## Family NOTONECTIDAE

*Notonecta spinosa* Hungerford

A Cordilleran species, so far in Canada, recorded only from British Columbia (Scudder 1977; Maw *et al.* 2000). The species can be keyed using Hungerford (1933).

New records: **AB**: 1 specimen, Waterton Lakes Nat. Park, grassland pond with mudstone, substrate, near Buffalo Paddock, 49°07'44"N 113°51'11"W, 9.vii.2005 (R.E. Roughley & R.D. Kenner) [UBC]; 1 specimen, Waterton Lakes Nat. Park, spring-fed *Carex* marsh, near West Entrance on Hwy. 5, 49°07'12"N 113°50'53"W, 9.vii.2005 (R.E. Roughley & R.D. Kenner) [UBC].

## Family PLEIDAE

*Neoplea striola* (Fieber)

This pygmy backswimmer, so far in Canada is recorded only from Manitoba, Ontario and Quebec (Maw *et al.* 2000). The species is keyed in Brooks and Kelton (1967).

New record. **BC**: 1 specimen, Edge-wood, F11, edge forest by beaver pond, 25.viii.-5.ix.1988 (H. Knight) [CNC].

## Infraorder GERROMORPHA

## Family VELIIDAE

*Microvelia pulchella* Westwood

This cosmopolitan species is recorded from Alaska, and in Canada from most provinces from British Columbia to Newfoundland (Maw *et al.* 2000). The species is keyed in Brooks and Kelton (1967).

New record. **SK**: 1♂ 2♀, Cowan Dam at Hwy. 55, 54°11'49"N 107°27'0"W, *Typha* pond, 22.vii.2003 (R.E. Roughley) [CNC].

## Family GERRIDAE

*Gerris incognitus* Drake & Hottes

A species reported across Canada from British Columbia to Labrador (Maw *et al.* 2000), but not previously recorded from Prince Edward Island. The species is keyed by Drake and Harris (1934).

New record. **PE**: 1♂ 1♀, Millburn, 8.vi.1970 (Ray Wenn) [UPEI].

*G. pingreensis* Drake & Hottes

This species is known from Alaska and across Canada from Yukon to Labrador (Maw *et al.* 2000), but to date there are no published records for New Brunswick, Prince Edward Island, Nova Scotia and Newfoundland. The species is keyed by Drake and Harris (1934) and Brooks and Kelton (1967).

New record. **NF**: 2♂ 1♀, Plum Point, ponds, 11.ix.1999 (T. Huxley) [CNC].

*Limnoporus notabilis* (Drake & Hottes)

This western water strider in Canada is known from British Columbia and Alberta (Maw *et al.* 2000; Andersen and Spence 1992). The species is keyed by Andersen and Spence (1992).

New record. **SK**: 1♂, Cypress Hills, 15.vii.1977 (R. Hopper) [SM].

## Family SALDIDAE

*Micracanthia humilis* (Say)

A Nearctic-Neotropical species, widely distributed in North America, and in Canada reported from British Columbia and Northwest Territories east to Newfoundland, but not previously recorded from Nova Scotia (Maw *et al.* 2000). The species

is keyed by Schuh (1967) and Polhemus and Chapman (1979).

New records. **NS:** 2♀, Halifax Co., Lawrencetown Beach, viii.1979 (B. Wright) [NSM]; 1♀ Halifax Co., Petpeswick Harbour, wet sandy upper shore, 29.vii.1971 (Barry Wright) [NSM].

#### *Salda lugubris* (Say)

A widely distributed Nearctic species that also occurs in Mexico, and in Canada is reported from Yukon to Newfoundland, but not previously from New Brunswick and Nova Scotia (Maw *et al.* 2000). The species is keyed by Brooks and Kelton (1967) and Schuh (1967).

New records. **NB:** 1♂, French Lake, 2.vii.1928 (W.S. Brown) [CNC]. **NS:** 1♀, CBI, Chiticamp, vi.-vii.1917 (F. Johansen) [CNC]; 2♀, Halifax Co., Port Wallis, 17.viii.1952 (D.C. Ferguson) [NSM].

#### *Saldula ablusa* Drake & Hottes

This is an eastern Nearctic species, keyed and reported from Ontario by Schuh (1967). Polhemus (1988) noted that references to *Saldula xanthochila* (Fieber) from the northeastern United States almost certainly refer to *S. ablusa* or *S. pallipes* (Fabricius). Wright (1989) reported *S. xanthochila* from Sable Is., Nova Scotia, and examination of 2♂ 1♀ specimens in the Nova Scotia Museum of Natural History, labeled "NS. Sable Is., brackish ponds east of station, 26.vii.1976 (Barry Wright)" shows these to be *S. ablusa*. As pointed out by Polhemus (1985), *S. ablusa* lacks a distinct dark distal streak on the ventral side of the hind femora, and as noted by Schuh (1967) typically has pale lateral margins to the pronotum, whereas these are never present in the *Saldula* "*pallipes* – *palustris*" group.

New record. **PE:** 7♂ 1♀, Can. Nat. Park, Dalvay House, 19.vii.1940 (G.S. Walley) [CNC].

#### *S. bouchervillei* (Provancher)

This species was previously reported from Nova Scotia as *Salda bouchervillei* (Provancher) by Wright (1989).

Material examined: **NS:** 2♀, Sable Island, brackish ponds east of station, 26.vii.1976 (Barry Wright) [NSM].

#### *S. laticollis* (Reuter)

Lindskog (1981) clarified the identity of this species, distinguishing it from the closely related *S. pallipes* (Fabricius) and *S. palustris* (Douglas) by the presence of long, curved, semi-recumbent or suberect setae on the head dorsally, and noted that *S. fernaldi* Drake is a synonym. In Old World populations of both *S. pallipes* and *S. palustris*, the pubescence on the head and dorsum generally is uniformly short and recumbent. However, as observed by Schuh (1967), the *Saldula* "*pallipes* – *palustris*" species complex has confused systematic heteropterists for some time, because of the extreme variability of the "species". Nevertheless, coastal populations from western North America and Newfoundland, previously identified as *S. palustris* have been shown to be *S. laticollis* (Lindskog 1981; Polhemus 1988), the intertidal biology of which has been described by Stock and Lattin (1976) under *S. palustris*.

Wright (1989) reported *S. palustris* from Sable Is., Nova Scotia, but examination of specimens in the Nova Scotia Museum of Natural History shows these to be *S. laticollis*. This species is also now known from New Brunswick and Prince Edward Island.

New records. **NB:** 1♂ 1♀, Kouchibouguac Nat. Park, 14.vi.1977 (S.J. Miller) [CNC]. **NS:** 2♂ 3♀, Digby Co., Sandy Cove, 4.viii.1971 (Barry Wright) [NSM]; 2♀, Guysborough Co., Liscombe, marshy area above beach, 13.viii.1971 (Barry Wright) [CNC]; 2♂ 1♀, Sable Island, West Light, 23.vii.1976 (Barry Wright) [NSM]; 1♂ 1♀, Sable Island, freshwater pond at West Light, 13.vi.1977 (Barry Wright) [NSM]. **PE:** 1♂, Brackley Beach, 5.vii.1966 (L.A. Kelton) [CNC]; 3♂, Green Gables, Cavendish Beach, 22.vii.1967 (J.E.H. Martin) [CNC].

### Infraorder CIMICOMORPHA

#### Family ANTHOCORIDAE

*Anthocoris tomentosus* Péricart

This western Nearctic, and Beringian species occurs in the western United States south to Arizona (Lewis *et al.* 2005), and Alaska to Manitoba, but has not previously been reported from Saskatchewan (Maw *et al.* 2000). The species is keyed in Kelton (1978) under the name *A. melanocerus* Reuter, and characteristically has the hemelytra entirely shiny, and the pronotum and antennae completely black. Lewis *et al.* (2005) reported *A. tomentosus* being regularly collected from *Alnus*, *Populus*, *Pyrus* and *Salix* growing in and near fruit-growing regions in Washington State, as well as on *Rumex* and psyllid-infested *Shepherdia argentea* (Pursh) Nutt.

New record. **SK**: 1♀, Fort Qu'Appelle, 17.vi.1967 (R. Hooper) [SM]; 1♀, Regina, on elm, 6.x.1986 (K. Roney) [SM].

*Dufouriellus ater* (Dufour)

This species was described from Europe, and was first reported from America north of Mexico by Van Duzee (1916). It is now known to be widely distributed in North America, with published records in the USA for New York (Van Duzee 1917), North Carolina (Blatchley 1926), California and Kentucky (Blatchley 1928), Idaho (Harris and Shull 1944), Oregon (Lattin 2004) and Hawaii (Lattin 2005, 2007a), whereas in Canada it has previously been reported from British Columbia (Anderson 1962) and Ontario (Kelton 1978).

*Dufouriellus ater* is usually collected under the bark of trees, but also is often associated with stored products (Awadallah *et al.* 1984; Arbogast 1984; Lattin 1999). As a result, it is a useful predator of some economic importance (Lattin 2000).

The species is keyed by Kelton (1978), who noted that in Canada it is rare in collections, and probably introduced into British Columbia and Ontario. Although *D. ater* was not included as an alien in Canada by Scudder and Foottit (2006), Lattin (2004, 2007b) considered the species to be non-indigenous to America north of Mexico.

New record. **NS**: 1♂, Halifax, Grain

Elevators, from dust and debris samples, 17.vii.1991 (J. Hulton) [NSM].

## Family NABIDAE

*Nabis inscriptus* (Kirby)

This Holarctic species, with somewhat abbreviated wings is difficult to distinguish from shorter winged specimens of *N. americanoferus* Carayon. Reliable separation is based on the shape of the male parameres and structure of the copulatory pouch of the female, as illustrated in Vinokurov (1988) and discussed by Kerzhner (1963). *Nabis inscriptus* is recorded from Alaska and across northern Canada (Maw *et al.* 2000), but as noted by Henry and Lattin (1988), reported distribution records need to be verified.

New records. **NS**: 1♀, Antigonish Co., Pomquet Beach, 31.v.1978 (B. Wright) [NSM]; 1♂, Halifax, found on corpse, Path. Lab., 1.xi.1979 (B. Wright) [NSM].

*N. roseipennis* Reuter

This species is recorded from British Columbia to Nova Scotia in Canada (Maw *et al.* 2000), but has not previously been reported from Prince Edward Island. The species is keyed in Blatchley (1926) and Harris (1928) and characteristically has black spots on the hind tibiae.

New record: **PE**: 6♂ 8♀, Charlottetown, UPEI, nr. East edge, 46°15'25"N 63°08'08"W, sweeping, 20.ix.2004 (G.G.E. Scudder) [CNC, UPEI].

*Pagasa nigripes* Harris

Kerzhner (1993) raised *Pagasa fusca* var. *nigripes* Harris to specific rank, and recorded the species in Canada from Alberta, Quebec and Saskatchewan. He also reported *P. nigripes* from Alaska, Colorado, Massachusetts, New Hampshire, New Mexico, New York, Pennsylvania, Vermont and Wyoming. He distinguished it from *Pagasa fusca* (Stein) by differences in the male and female genitalia, and noted that the legs tend to be brown or brownish yellow in *P. nigripes*, whereas they are yellow in *P. fusca*, with the femora often orange or reddish. However, the legs in *P. fusca* can

sometimes be partly brownish or entirely black. Hence, the genitalic characters are the most reliable. The parameres in *P. fusca* are relatively large with the outer margin rounded, whereas in *P. nigripes* they are slightly smaller than in *P. fusca* and distinctly angulate on the outer margin.

*Pagasa nigripes* is here recorded for the first time from British Columbia, Northwest Territory and Yukon, where the species was previously reported as *P. fusca* in Maw *et al.* (2000): *P. fusca* does not occur in Alaska, Northwest Territory and Yukon. However, in British Columbia, *P. fusca* co-occurs with *P. nigripes*, the species having been collected together at Merritt (23 km E, Hamilton Commonage), Osoyoos (Mt. Kobau in Montane Spruce habitat), Vaseux Creek (Kennedy bench), and the Windermere Valley.

New records. **BC**: 1♂, Canal Flats, 10.8 km S., 31.viii.1998 (G.G.E. Scudder) [CNC]; 1♂, Fairview, White L., BGxh1, SWm, pan trap WL/P-1, 4.vii.-11.vii.1995 (J. Jarrett) [UBC]; 3♀, Merritt, 23 km E, Hamilton Commonage, Upper Fescue grassland, early seral, 1250 m, 16.ix.2000 (G.G.E. Scudder) [CNC, UBC]; 1♀, Merritt, 35 km S, 14.viii.1988 (G.G.E. Scudder) [CNC]; 1♂, Nicola, 24.vii.1932 (G.J. Spencer) [UBC]; 1♂, Osoyoos, East Bench, *Artemisia/Purshia* assoc., pitfall trap, 15.vii. 17.viii. 1990 (G.G.E. Scudder) [CNC]; 1♂, Osoyoos IRI, Inkaneep, BGxh1, AN, pitfall trap T1-1, 6.vii.-9.viii.1995 (G.G.E. Scudder) [CNC]; 1♂, *id.*, 9.viii.-9.ix.1995 [CNC]; 1♀, *id.*, 9.ix.-6.x.1995 [CNC]; 1♀, *id.*, T2-1, 9.viii.-9.ix.1995 [CNC]; 1♂, *id.*, T2-5, 4.vii.-7.ix.1994 [CNC]; 1♂, *id.*, T4-1, 9.viii.-9.ix.1995 [CNC]; 1♀, *id.*, T4-3, 6.vii.-9.viii.1995 [CNC]; 1♀, *id.*, T4-5, 4.viii.-7.ix.1994 [CNC]; 1♀, *id.*, T5-2, 9.viii.-9.ix.1995 [CNC]; 1♂, Osoyoos, Mt. Kobau, MSxh, VK, Pitfall trap K4A-5, 10.viii.-8.ix.1995 (J. Jarrett) [UBC]; 1♀, *id.*, K4A-3, 18.viii.-28.ix.1997 [UBC]; 1♂, Osoyoos, Mt. Kobau Rd., IDFdk1, pitfall trap K3B-5, 18.viii.-28.ix.1996 (J. Jarrett) [UBC]; 1♂, *id.*, IDFxh1, pitfall trap K2B-4, 18.viii.-28.ix.1996 [UBC]; 1♂, *id.*, PPxh1, pitfall

trap K1A-2, 28.vii.-18.viii.1997 [UBC]; 1♂ 1♀, Tatlayoka L., 16.vii.1978 (G.G.E. Scudder) [CNC]; 1♀, Vaseux Cr., CWS bench, BGxh1, AN, pitfall trap Y1-4, 6.ix.-4.x.1995 (G.G.E. Scudder) [CNC]; 1♀, *id.*, Y2-5 [CNC]; 1♂, *id.*, Y1-4, 8.vii.-3.viii.1994 [CNC]; 1♀, *id.*, Y2-5, 12.viii.-6.ix.1995 [CNC]; 1♀, *id.*, Y2-5, 6.ix.-4.x.1995 [CNC]; 1♀, *id.*, Y3-3, 5.vii.-12.viii.1995 [CNC]; 2♂, *id.*, Y3-3, 6.ix.-4.x.1995 [CNC]; 1♂, *id.*, Y3-5, 5.vii.-12.viii.1995 [CNC]; 1♂, *id.*, Y4-1, 12.viii.6.ix.1995 [CNC]; 1♂, *id.*, Y4-5, 3.viii.-6.ix.1994 [CNC]; 1♂, *id.*, Y4-5, 5.vii.-12.viii.1995 [CNC]; 1♂, Vaseux Cr., 'Kennedy bench', 49°16'N 119°30'W, BGxh1, AN, pitfall trap Z2-4, 3.vi.-8.vii.1994 (G.G.E. Scudder) [CNC]; 1♀, *id.*, Z2-5, 12.viii.-6.ix.1995 [CNC]; 1♂, Vaseux Cr., 'Kennedy flats', 49°15'N 119°31'W, BGxh1, AN, pitfall trap X2-1, 6.ix.-4.x.1995 (G.G.E. Scudder) [CNC]; 1♀, *id.*, X3-4 [CNC]; 1♂, X3-5, 12.vii.-6.ix.1995 [CNC]; 1♀, *id.*, X4-1 [CNC]; 1♂, Vaseux L., Wildlife Res., BGxh1, AN:F, pitfall trap VL1-2, 27.vii.-17.viii.1997 (J. Jarrett) [UBC]; 1♂, Westwick L., Cariboo, 28.vi.1961 (J. Scudder) [UBC]; 1♀, *id.*, 17.viii.1962 (G.G.E. Scudder) [CNC]; 3♂ 4♀, *id.*, 18.viii.1962 [CNC, RBCM, UBC]; 1♀, White L., BGxh1, SWm, pitfall trap WL2-3, 17.viii.-28.ix.1996 (J. Jarrett) [UBC]; 1♂ 2♀, Windermere Valley, pitfall trap No. 5, 26.vii.-17.ix.2000 (R. Sargent) [CNC, UBC]. **NT**: 1♀, Fort Smith, 27.v.1950 (W.G. Helps) [CNC]; 1♀, Fort Smith, 6.viii.1950 (J.B. Wallis) [CNC]. **YT**: 2♀, Alaska Hwy. km 1768, Duke R., 9.vii.1983 (G.G.E. Scudder) [CNC]; 1♂, Alaska Hwy. mi 1054, Kluane L., 16.vii.1962 (G.G.E. Scudder) [CNC]; 1♂, *id.*, 7.vii.1983 [CNC]; 2♂, Canyon, Aishihik R., 9.vii.1983 (G.G.E. Scudder) [CNC]; 3♂ 3♀, Carcross, 8.vii.1983 (G.G.E. Scudder) [CNC, UBC]; 1♂, Lapie R., 1 km E on Campbell Hwy., 28.vii.1981 (C.S. Guppy) [UBC]; 1♀, Pelly Crossing, 17.vii.1983 (G.G.E. Scudder) [CNC]; 1♀, Tatchun Cr., 62°17'N 136°17'W, 17.vii.1983 (G.G.E. Scudder) [CNC]; 1♀, Whitehorse, 17.vii.1959 (R. Madge) [CNC]; 1♀, White-

horse, 31.vii.1981 (C.S. Guppy) [UBC].

I have also collected *P. nigripes* in Washington State, USA, as follows: WA: 1♀, Oroville, E. Osoyoos L., 48°53'N 119°25'W, *Purshia* assoc., AN BGxh1, pitfall trap O2-1, 10.ix.-4.x.1995 (G.G.E. Scudder) [CNC].

#### Family MIRIDAE

##### *Ceratocapsus modestus* (Uhler)

A widely distributed eastern Nearctic species, previously reported from Saskatchewan east to Quebec in Canada (Maw *et al.* 2000). Recorded hosts in West Virginia are *Quercus alba* L. and *Vitis* sp. (Wheeler *et al.* 1983). The species is keyed by Knight (1941), Henry (1979), Kelton (1980) and Larochelle (1984).

New records. NS: 1♀, Grand Pre, on *Picea*, 10.viii.1966 (L.A. Kelton) [CNC]; 1♀, Grand Pre, *Pinus sylvestris*, 10.viii.1966 (L.A. Kelton) [CNC].

##### *Conostethus americanus* Knight

To date this species in Canada has been reported only from Alberta, Northwest Territories and Saskatchewan (Maw *et al.* 2000). On the prairies *C. americanus* occurs on grasses (Kelton 1980). It is keyed and illustrated in Kelton (1980).

New record. YT: 4♂ 7♀, Whitehorse, Dillabough's graze lease, 8V 6754911 490889, 12.vii.2005 (G.E. Hutchings) [RBCM].

##### *Cyrtorhinus caricis* (Fallén)

A Holarctic species, reported to occur on sedge (*Carex* spp.) across Canada (Kelton 1980), and also recorded from Alaska and Minnesota, with an apparent relict population occurring in Colorado (Wheeler and Henry 1992). The species is keyed and illustrated by Kelton (1980).

New record. NS: 1♂, Lake Egmont, 18.vii.1991 (B. Wright) [NSM].

##### *Deraeocoris quercicola* Knight

An eastern Nearctic species, widely distributed and in Canada reported to date from Saskatchewan east to Quebec (Maw *et al.* 2000). Recorded hosts include *Carya*

sp., *Quercus alba*, *Q. ilicifolia* Wangenh and *Tilia americana* L. (Wheeler *et al.* 1983), as well as *Quercus macrocarpa* Michx. where it preys on aphids (Kelton 1980). The species is keyed by Knight (1921), Kelton (1980) and Larochelle (1984).

New record. NB: 1♀, St. Johns, Rockwood Pk., 5.viii.1954 (J.F. Brimley) [CNC].

##### *D. triannulipes* Knight

This Nearctic species in Canada has previously been reported from British Columbia east to Quebec (Maw *et al.* 2000). *Deraeocoris triannulipes* is reported to feed on aphids on *Populus tremuloides* Michx. and *Alnus* spp. (Kelton 1980). It is keyed by Knight (1921), Kelton (1980) and Larochelle (1984).

New records. NS: 1♂, Exfern, on apple, 4.vii.1950 (F.T. Low) [CNC]; 1♀, Halifax, *Pyrus*, 22.vii.1976 (L.A. Kelton) [CNC]; 2♀, Kentville, on apple, 10-14.vii.1976 (L.A. Kelton) [CNC]; 1♂, Kentville, *Tilia cordata*, 15-17.vii.1976 (L.A. Kelton) [CNC].

##### *Dichaetocoris piceicola* (Knight)

A western Nearctic species, known from Colorado in the USA, and in Canada recorded from Alberta, British Columbia and Yukon (Maw *et al.* 2000). The genus is keyed by Knight (1968). The species *D. piceicola* is distinguished from the only other northern species (*D. gillespiei* Schwartz and Scudder) by Schwartz and Scudder (2003). It is recorded from *Picea engelmanni* Parry (Polhemus 1994), and in British Columbia has been collected on *Abies lasiocarpa* (Hook.) Nutt., *Picea* sp., *Pinus contorta* Dougl. and *Tsuga heterophylla* (Raf.) Sarg. (Scudder, unpublished).

New record. AK: 1♀, Mosquito L., 59°27'N 136°02'W, 6.vii.1983 (G.G.E. Scudder) [CNC].

##### *Labops verae* Knight

A western Nearctic and Beringian species distributed from Alaska to Manitoba and south to Washington State (Henry and

Wheeler 1988; Maw *et al.* 2000), but not previously reported from Saskatchewan. The species is keyed in Slater (1954) and Kelton (1980). The host plants are unknown (Kelton 1980).

New record. **SK**: 1♂, Stony Rapids, 30.vi.1975 (R. Hooper) [SK].

#### *Lygidea salicis* Knight

This Nearctic species in Canada is previously reported from Alberta east to Newfoundland (Maw *et al.* 2000), and in the United States from New York to Minnesota, Colorado, and California (Henry and Wheeler 1988). It has not previously been recorded from British Columbia. *Lygidea salicis* Knight is a small species with the average length in the male of 5.8 mm, and in the female 6.2 mm. The species is keyed by Kelton (1980), who notes that the pilosity on the second antennal segment is shorter than the thickness of this segment. *Lygidea salicis* is usually collected on *Salix* spp. (Kelton 1980; Wheeler *et al.* 1983).

New records. **BC**: 2♂ 3♀, Fernie, gold-enrod, 23.vii.1959 (L.A. Kelton) [CNC]; 2♂ 2♀, Mt. Revelstoke Nat. Pk., *Salix*, 17.vii.1970 (L.A. Kelton) [CNC].

#### *Megalopsallus femoralis* Kelton

This species so far has been reported from Alberta, Manitoba, Saskatchewan, Colorado, South Dakota and Wyoming (Schuh 2000b). *Megalopsallus femoralis* has been collected on *Salicornia rubra* Nels. (Kelton 1980). It is keyed by Kelton (1980) and Schuh (2000b), and is illustrated in colour in the latter reference.

New record. **BC**: 3♂ 2♀, Kamloops, Ironmask L., 10U 6804 56152, saline flats, *Salicornia/Plantago*, 730 m., 14.vi.1995 (S.G. Cannings) [RBCM].

#### *Orthotylus alni* Knight

This Nearctic species is distributed from Yukon to Newfoundland, and south to New York and Minnesota in the eastern United States (Henry and Wheeler 1988; Maw *et al.* 2000). However, it has not previously been recorded from Nova Scotia. *Orthotylus alni* is keyed by Kelton (1980), and has

been collected on *Alnus rugosa* (DuRoi) Spreng. (Kelton 1980), as well as *A. tenuifolia* Nutt., *Betula glandulosa* Michx., *Lupinus* sp. and *Salix* sp. (Scudder 1997).

New records. **NS**: 1♂, Chester, 10.vii.1969 (B. Wright) [NSM]; 1♀, Chester, 16.vii.1969 (B. Wright) [NSM].

#### *O. nyctalis* Knight

Described originally from Minnesota (Knight 1927), this species has been reported in the USA also from Iowa, Illinois, New York and Wisconsin (Henry and Wheeler 1988). It has not previously been recorded from Canada under this name.

According to Knight (1927), *O. nyctalis* can be recognized chiefly on the structure of the male genital claspers. The left clasper is slender with two short dorsal prongs, and the right clasper decurved on the apex and devoid of spines, but the dorsal margin has a prominent spine at the basal third, and two other spines just before the decurved apex.

The record of *O. candidatus* Van Duzee from Saskatchewan (Kelton 1980; Maw *et al.* 2000; Roch 2007) is evidently referable to *O. nyctalis*, as is the record of *O. candidatus* from Ontario (Maw *et al.* 2000; Roch 2007). The recorded occurrence of *O. candidatus* in Quebec (Henry and Wheeler 1988; Roch 2007) may also refer to *O. nyctalis*. Some specimens from Dawson and Moose Creek in the Yukon, listed as *Orthotylus* sp. in Scudder (1997) are actually *O. nyctalis* (see below), but it may be noted that *O. candidatus* also occurs at both these localities. *Orthotylus nyctalis* evidently occurs on *Populus tremuloides* (Kelton 1980).

New records. **AB**: 1♂, Stettler, 3.viii.1957 (A. & J. Brooks) [CNC]; 1♀, Vermilion Provincial Park, Beaverdam Loop Trail, *Populus tremuloides* Michx., 22.viii.1993 (M.D. Schwartz) [CNC]. **MB**: 1♂, Falcon L., 5.viii.-10.viii.1978 (L.A. Kelton) [CNC]; 1♀, *id.*, 6.viii.1978 (L.A. Kelton) [CNC]; 1♀, Rennie, 16.viii.1961 (F.I.S.) [CNC]. **ON**: 1♂, One Sided Lake, *Salix* sp., 1.viii.1960 (Kelton and Whitney) [CNC]; 1♀, Tillsonburg, 18.vii.1962

(Kelton and Thorpe) [CNC]. **SK**: 1♀, Cypress Hills Prov. Park., *Sheperdia canadensis*, 19.ix.1951 (L.A. Konotopetz) [CNC]. **YT**: 1♂, Dawson, 14 mi E, *Populus* sp., 29.vii.1962 (R.E. Leech) [CNC]; 1♀, Dawson, *Salix* sp., 23.vii.1983 (L.A. Kelton) [CNC]; 1♀, Moose Creek, *Salix* sp., 28.vii.1983 (L.A. Kelton) [CNC].

*Phytocoris buenoi* Knight

An eastern Nearctic species, in Canada previously reported from Ontario and Quebec (Maw *et al.* 2000). The species is keyed by Blatchley (1926). Knight (1920) reported it to occur on Norway spruce (*Picea abies* (L.) Karst.) in the eastern United States, and Wheeler *et al.* (1983) added *Picea glauca* (Moench) Voss and *P. rubens* Sarg.

New records. **NS**: 1♂, Chester, 29.viii.1968 (B. Wright) [NSM]; 1♀, Sandy Cove, 4.viii.1971 (B. Wright) [NSM]; 1♀, Baddeck, 28.viii.1972 (B. Wright) [NSM]; 1♂, Kemptville, 24.viii.1982 (Agriculture Canada) [NSM].

*P. procteri* Knight

This species, which is a member of the *P. junceus* Knight group, was described from Maine (Knight 1974), and is reported from Quebec (Roch 2007). The frons has definite transverse red lines, the pronotum is pallid with a basal submarginal strong black band, and the propleura are brownish black. The clavus is more or less fuscous external to the claval vein, with insect length of 8.0 mm. The first antennal segment is pallid, but clothed with black, recumbent setae, and without distinct dark spots. The first antennal segment is longer than the width of the vertex, but does not exceed the width of the pronotum. The second antennal segment is without annuli or coloured bands. There are no records of a host for this species.

New records. **NS**: 1 specimen (abdomen missing), Lake Kejimikujik, 13.vii.1961 (D.C. Ferguson) [NSM]; 1♂, Chester, 24.vii.1968 (B. Wright) [NSM].

*Pilophorus neoclavatus* Schuh & Schwartz

This eastern Nearctic species to date in Canada has been reported from Alberta east to Quebec (Maw *et al.* 2000), and is keyed by Schuh and Schwartz (1988). Hosts include *Alnus rugosa*, *Quercus ilicifolia*, *Q. palustris*, *Q. stellata* Wangenh and *Salix longifolia* Muhl. (Schuh and Schwartz 1988).

New record. **NS**: 1♀, Kentville, 8.viii.1952 (C.R. McL.) [LM].

*Rhinocapsus rubricans* (Provancher)

An eastern Nearctic species, in Canada to date reported from Saskatchewan east to Quebec (Maw *et al.* 2000). The species is keyed and illustrated in Kelton (1980), who reports it collected on *Kalmia polifolia* Wang.

New record. **NS**: 1♂, Chester, 4.vii.1969 (B. Wright) [NSM].

*Sixeonotus deflatus* Knight

An eastern Nearctic species, in Canada previously only reported from Quebec (Larochelle 1984; Maw *et al.* 2000). The species is keyed by Larochelle (1984) as *S. insignis* Reuter, who records the host as *Symplocarpus foetidus* (L.).

New records. **NB**: 1♂, St. John, 9.viii.1954 (J. Brimley) [CNC]. **ON**: 8♂ 5♀, St. Catherines, 22.vi.1961 (Kelton and Brampton) [CNC].

Family TINGIDAE

*Acalypta lillianis* Torre-Bueno

This Nearctic tingid is widely distributed in North America and Beringia (Drake and Lattin 1963; Scudder 1997). It occurs in Alaska, and from Yukon to Newfoundland, but has not previously been reported from Manitoba (Maw *et al.* 2000). It is keyed by Drake and Lattin (1963), who note that host records are mosses.

New record. **MB**: 1♂, Bird Cove, 4 km NE, Churchill North Studies Centre, 58° 46'14"N 93°50'33"W, pit trap in tundra zone, 18.viii.2006 (Boreal & Arctic Entomol.) [UM].



## Family ARADIDAE

*Aradus uniannulatus* Parshley

A Nearctic and Beringian species, distributed from Yukon to Quebec, and in the United States, south to New York and Colorado (Scudder 1997). The species is keyed by Matsuda (1977) who notes that it is reported to be associated with *Pinus contorta murrayana* (Balf.) Critchfield elsewhere.

New records. **BC**: 1♀, Fernie, 22.viii.1934 (Hugh Leech) [CNC]; 1♀, Lorna, *Picea engelmanni*, 17203 Lot 1, 29.vi.1929 (R. Hopping) [CNC]; 1♂, *id.*, 17203 Lot 2, 10.vii.1924 [CNC]; 1♀, *id.*, *Pinus contorta*, 17203 Lot 8, 12.vii.1924 [CNC]; 1♂, *id.*, *Picea engelmanni*, 17203 Lot 14, 12.vii.1924 [CNC]; ♂, *id.*, 17203 Lot 25, 25.vii.1924 [CNC]; 1♂, *id.*, 17203 Lot 26, 27.vii.1924 [CNC] 1♀, Midday Valley, Merritt, *Pinus ponderosa*, Exp. 17501 Lot 1035, 30.v.1923 (R. Hopping) [CNC]; 1♀, Pine Pass, *Picea*, 11.vii.1972 (D.E. Bright) [CNC].

## Infraorder PENTATOMOMORPHA

## Family COREIDAE

*Leptoglossus occidentalis* Heidemann

This leaf-footed bug, commonly called the western conifer seed bug, feeds on numerous conifer species (Koeber 1963; Krugman 1969; Hedlin *et al.* 1981; Schaefer and Mitchell 1983; Gall 1992; Mitchell 2000) although the apparently strong reliance on Pinaceae as a food source is not absolute (Mitchell 2000). However, it is of considerable economic importance because it can cause significant losses in conifer seed orchards (Koeber 1963; Schowalter and Sexton 1990; Blatt and Borden 1996; Mitchell 2000; Strong *et al.* 2001; Bates *et al.* 2002; Strong 2006). The species is keyed in Allen (1969) and McPherson *et al.* (1990), and illustrated by Koeber (1963) and Ruth *et al.* (1982).

Originally considered a western Nearctic species, *L. occidentalis* has naturally expanded its range eastwards in the past few decades (Schaffner 1967; McPherson *et al.* 1990; Marshall 1991; Gall 1992; Ridge-O'Connor 2001), and has invaded Europe

(Taylor *et al.* 2001; Gogala 2003; Tescari 2004; Hilpold 2005; Rabitsch and Heiss 2005; Ribes and Oleguer 2005; Foldessy 2006; Moulet 2006). It is now known to occur in Nova Scotia.

New records. **NS**: 1♀, Kings Co., Middleton, in house, 20.ix.2006 (J. Parks) [NSM]; 1♀, Halifax Co., Halifax, on house, 3.x.2006 (B. Fay) [NSM]; 2♀, Kings Co., Lakeville, 1 of 4, 12.iii.2007 (J. Morton) [NSM 36185]; 1♀, Halifax Co., Halifax, in dwelling, 15.x.2007 (John Sherwood) [NSM Cat. 36219].

## Family RHOPALIDAE

*Liorhyssus hyalinus* (Fabricius)

This cosmopolitan species is widely distributed in North America, and in Canada has so far been reported from British Columbia, Manitoba and Ontario (Maw *et al.* 2000). It is keyed by Blatchley (1926), Slater and Baranowski (1978), and Hoebeke and Wheeler (1982), and illustrated in Slater and Baranowski (1978). Schaefer and Chopra (1982) report *Abutilon*, *Euphorbia*, *Lactuca* and *Sonchus* as host plant genera.

New record. **QC**: 1♀, Terrebonne Co., Lac Carre, Lot 31, Range 8, 19-23.viii.1968 (W. Boyle and R. La Conde) [LM].

## Family LYGAEIDAE

*Kleidocerys ovalis* Barber

This widely distributed Nearctic species has so far in Canada only been recorded from British Columbia and Ontario (Maw *et al.* 2000), as well as Quebec (Roch 2007). It is keyed in Barber (1953) and Scudder (1962). In British Columbia, *K. ovalis* has been collected on *Abies lasiocarpa*, *Alnus* sp., *Betula occidentalis* Hook., *B. papyrifera* Marsh., *Malus* sp., and *Pinus ponderosa* Dougl. (Scudder, unpublished).

New records. **AB**: 5♂ 2♀, Drumheller, 18.vi.1957 (Brooks, MacNay) [CNC]; 7♂ 6♀, *id.*, 11.viii.1957 (A.R. & J.E. Brooks) [CNC]; 2♂ 3♀, Empress, 7.vi.1957 (Brooks, MacNay) [CNC]; 1♀, Lethbridge, 3.vii.1929 (J.H. Pepper) [CNC]; 1♂, Lundbreck, 7.viii.1930 (J.H. Pepper) [CNC]. **MB**: 1♀, Aweme, 5.vii.1920 (H.A. Robertson) [CNC]; 1♀, *id.*, 8.vii.1920 (P.N.

Vroom) [CNC]; 3♀, *id.*, *Betula*, 30.vi.1922 (N. Criddle) [CNC]; 1♀, *id.*, 3.v.1923 (N. Criddle) [CNC]; 1♂ 1♀, Carberry, 9.v.1953 (Brooks, Kelton) [CNC]; 1♂ 3♀, Ninette, *Betula glandulosa*, 21.vi.1958 (J.F. McAlpine) [CNC]; 1♂, *id.*, 14.vii.1958 (R.B. Madge) [CNC]; 1♂, *id.*, 15.vii.1958 (R.L. Hurley) [CNC]; 1♂, Onah, 10.v.1923 (R.M. White) [CNC]; 8♂ 1♀, *id.*, *Betula papyrifera*, 10.ix.1930 (R.M. White) [CNC]; 1♀, Turtle Mtn., 22.vii.1953 (Brooks, Kelton) [CNC]. **SK**: 1♀, Punnichy, 21.v.1965 (R. Hooper) [SM].

#### *K. resedae* (Panzer)

This Holarctic species is widely distributed in North America, and occurs in Alaska and from Yukon to Newfoundland and Labrador, but has not previously been recorded from Prince Edward Island. It is keyed in Barber (1953) and Scudder (1962). *Kleidocerys resedae* usually occurs on *Alnus* spp. and *Betula* spp. (Scudder 1997).

New records. **PE**: 3♂ 1♀, Blooming Point, 46°24'33"N 62°58'07"W, sweeping, 20.x.2004 (G.G.E. Scudder) [CNC, UPEI]; 2♂ 4♀, Charlottetown, UPEI, nr. NE point, 46°15'39"N 63°08'19"W, sweeping, 20.x.2004 (G.G.E. Scudder) [CNC, UPEI].

#### *Melacoryphus lateralis* (Dallas)

A widely distributed Nearctic species, so far only recorded from British Columbia and Saskatchewan in Canada (Maw *et al.* 2000). The species is keyed by Slater (1988). Specimens of *M. lateralis* collected at light in Wyoming contained cardenolides in the body (Scudder and Duffey 1972), and thus showed evidence of feeding on Asclepiadaceous host plants.

New record. **ON**: 1♂, Guelph, 3.viii.1977 (W.A. Attwater) [UG].

#### Family RHYPAROCHROMIDAE

##### *Antillocoris minutus* (Bergroth)

An eastern Nearctic species, in Canada previously recorded from Ontario east to Newfoundland (Maw *et al.* 2000). The genus is keyed in Blatchley (1926), Slater and Baranowski (1978), and Larochelle (1984), with key to species given by Barber (1952)

and Larochelle (1984). The biology of *A. minutus* in New England has been described by Sweet (1964), who notes the species typically occurs on the ground and usually is found in forest litter, most frequently found beneath gray birch (*Betula populifera* Marsh) and white birch (*B. papyrifera*), but also occurs under hemlock (*Tsuga canadensis* (L.) Carr.), and in sphagnum bogs.

New record. **MB**: 1♀, Winnipeg, St. Charles Rifle Rge., Block B Refuge, Pitfall trap, 6-13.x.1999 (D.A. Pollock, J.K. Diehls and R.E. Roughley) [UM].

##### *Drymus umus* (Say)

An eastern Nearctic species, in Canada so far recorded from Saskatchewan east to Nova Scotia (Maw *et al.* 2000). The species is keyed by Blatchley (1926) and Larochelle (1984), and illustrated by Slater and Baranowski (1978). Sweet (1964) described the biology *D. umus* in New England, and noted that this is a ground-dwelling species, most abundant in subclimax forests, particularly where black birch (*Betula lenta* L.) and red maple (*Acer rubrum* L.) are associated with oak (*Quercus* spp.) and hickory (*Carya* spp.).

New record. **NF**: 1♂, St. John's, Long Pond, ix.-x.2001 (Biology 4150) [MU].

##### *Perigenes constrictus* (Say)

This eastern Nearctic species is distributed throughout the northern and central United States, and in Canada so far reported from Nova Scotia, Ontario and Quebec (Maw *et al.* 2000). The species is keyed by Blatchley (1926) and Larochelle (1984), and illustrated by Blatchley (1926) and Slater and Baranowski (1978). Sweet (1964) described the biology of *P. constrictus* in New England and noted that it typically occurs in temporary habitats, such as vacant lots, roadsides and newly fallow fields.

New record. **SK**: 1♂, Big Beaver, 9.vii.1974 (R. Hooper) [SM].

##### *Plinthisus americanus* Van Duzee

In Canada previously reported from Alberta east to New Brunswick (Maw *et al.*

2000). *Plinthisus americanus* is distinguished by the hemelytra of the female being densely pilose as noted by Sweet (1964), whereas in the closely related *P. compactus* (Uhler) the hemelytra of the female are glabrous. Sweet (1964) noted that *P. americanus* is a forest species in New England, most abundant in *Tsuga* litter.

New record. **BC**: 1♀, Attachie, 32 km W of Charlie L., 5.viii.1982 (R.A. Cannings) [RBCM].

*Trapezonotus arenarius* (Linnaeus)

A Holarctic species with a wide distribution in both the Nearctic and Palearctic, in Canada recorded from British Columbia and Yukon east to Quebec (Maw *et al.* 2000), and in the United States evidently restricted to the highlands of New England and northern New York (Sweet 1964). The species is keyed by Blatchley (1926) and Laroche (1984), and illustrated by Slater and Baranowski (1978). Sweet (1964) reported that in New England *T. arenarius* is a species of open upland habitats, particularly well-drained and rather dry sites.

New record. **NS**: 1♂, Lunenburg, 7.viii.1991 (B. Wright) [NSM].

Family CYDNIDAE

*Amnestus pusillus* Uhler

A widely distributed species in North America, with recorded occurrence also in Mexico and Guatemala (Froeschner 1960). In Canada, so far reported only from Ontario and Quebec (Maw *et al.* 2000). The species is keyed in Froeschner (1960), McPherson (1982), and Laroche (1984). McPherson (1982) reported that elsewhere *A. pusillus* has been collected from vegetation along streams and margins of roadsides and cultivated fields, as well as beneath rubbish in sandy places.

New record. **NB**: 1♀, Woodstock, 22.v.1966 (L.A. Kelton) [CNC].

Family PENTATOMIDAE

*Acrosternum hilare* (Say)

A widely distributed Nearctic species, in Canada so far recorded only from British

Columbia, Ontario and Quebec (Maw *et al.* 2000). This species has been collected from numerous plants (McPherson 1982) and can damage some crops (Panizzi *et al.* 2000). The species is keyed in Blatchley (1926), McPherson (1982), Rolston (1983), and Laroche (1984).

New record. **NS**: 1♀, Debert, 1.ix.1952 (R.L. Horsburgh) [LM].

*Cosmopepla intergressus* (Uhler)

A widely western Nearctic species, in Canada so far recorded only from British Columbia (Maw *et al.* 2000). *Cosmopepla intergressus* is keyed by McDonald (1986), who reported records of the species on "currants", *Rubus parviflorus* Nutt. and *Ribes* sp.

New record. **AB**: 3♂ 3♀, Lethbridge, black current, 21.ix.2005 (J.R. Byers) [AAFCL].

*Euschistus servus euschistoides* (Vollenhoven)

A Nearctic species widely distributed in North America, and in Canada recorded from British Columbia to Nova Scotia, but not previously reported from New Brunswick (Maw *et al.* 2000). It has been recorded from numerous host plants (McPherson 1982), and the species has caused yield and quality losses to several crops (Panizzi *et al.* 2000). The species is keyed in McPherson (1982) and Laroche (1984).

New records. **NB**: 1♀, Jonah Mt., 3.vi.1976 (P. Kevan) [LM]; 1♀, Whittier Ridge, 30.v.1976 (P. Kevan) [LM].

*Meneles insertus* (Say)

Widely distributed in North America, and in Canada previously recorded from Nova Scotia, Ontario and Quebec. *Meneles insertus* is nocturnal, arboreal, and phytophagous, and has been collected on deciduous trees (McPherson 1982). The species is keyed in Blatchley (1926), Rolston (1973), McPherson (1982) and Laroche (1984).

New record. **SK**: 1♀, Buffalo Pound Park, 5.v.1975 [SM].

*Zicrona caerulea* (Linnaeus)

This Holarctic species is widely distributed in North America, Europe and Asia, and occurs in the Oriental region (De Clercq 2000). In Canada it is reported from British Columbia east to New Brunswick (Maw *et al.* 2000). This predaceous species attacks only small prey (De Clercq 2000), and is keyed by McPherson (1982), Laroche (1984), and Thomas (1992).

New record. **NF**: 1♂, Red Indian L., Winddrift Lot 5, 25.vi.1980 (Brennan and Larson) [MU].

## Family THYREOCORIDAE

*Corimelaena pulicaria* (Germar)

A widely distributed species in North America, also reported from Mexico and Guatemala. In Canada recorded from British Columbia to Nova Scotia, but not previously reported from New Brunswick (Maw *et al.* 2000). The species has been collected on many plants (McPherson 1982), and is keyed in Blatchley (1926), McPherson (1982), and Laroche (1984).

New record. **NB**: 1♀, Fredericton, French Lake, 10.vi.1931 (C.W. Maxwell) [LM].

## ACKNOWLEDGEMENTS

The research for this paper was supported by grants from the Natural Sciences and Engineering Research Council of Canada. I thank the curators of the various collections for permission to examine the material in their care and/or loan of specimens. Dr. M.D. Schwartz helped me in my study of the Canadian National Collection of Arthropods in Agriculture and Agri-

Foods Canada in Ottawa, kindly identified or confirmed the identity of most of the Miridae listed, and allowed me to include the records of *Orthotyphus nyctalis*. Dr. T.J. Henry (United States Department of Agriculture, Washington, DC) provided useful comments on the manuscript. I thank Launi Lucas for the final preparation of the paper.

## REFERENCES

- Allen, R.C. 1969. A revision of the genus *Leptoglossus* Guerin (Hemiptera: Coreidae). *Entomologica Americana* (New Series) 45:35-140.
- Andersen, N.M. and J.R. Spence. 1992. Classification and phylogeny of the Holarctic water strider genus *Limnoporus* Stål (Hemiptera, Gerridae). *Canadian Journal of Zoology* 70:753-785.
- Anderson, N.H. 1962. Anthocoridae of the Pacific Northwest with notes on distribution, life histories, and habits (Heteroptera). *Canadian Entomologist* 94:1325-1334.
- Arbogast, R.T. 1984. Demography of the predaceous bug *Dufouriellus ater* (Hemiptera: Anthocoridae). *Environmental Entomology* 13:990-994.
- Awadallah, K.T., F.S. Tawfik, N. Abou-Zeid and El-Husseini. 1984. The life-history of *Dufouriellus ater* (Duf.) (Hemiptera: Anthocoridae). *Bulletin de la Société Entomologique d'Égypte* 63:191-197.
- Barber, H.G. 1952. The genus *Antillocoris* Kirk. in the United States (Hemiptera-Lygaeidae). *Bulletin of the Brooklyn Entomological Society* 47:85-87.
- Barber, H.G. 1953. A revision of the genus *Kleidocerys* Stephens in the United States (Hemiptera: Lygaeidae). *Proceedings of the Entomological Society of Washington* 55:273-283.
- Barnes, D.I., H.E.L. Maw and G.G.E. Scudder. 2000. Early records of alien species of Heteroptera (Hemiptera: Prosorrhyncha) in Canada. *Journal of the Entomological Society of British Columbia* 97:95-102.
- Bates, S.L., C.G. Lait, J.H. Borden and A.R. Kermode. 2002. Measuring the impact of *Leptoglossus occidentalis* (Hemiptera: Coreidae) on seed production in lodgepole pine using an antibody-based assay. *Journal of Economic Entomology*. 95:770-777.
- Blatchley, W.S. 1926. *Heteroptera or True Bugs of Eastern North America*, with especial reference to the Faunas of Indiana and Florida. Nature Publishing Company, Indiana. 1116 pp.
- Blatchley, W.S. 1928. Notes on the Heteroptera of eastern North America with descriptions of new species. 1. *Journal of the New York Entomological Society* 36:1-23.
- Blatt, S.E. and J.H. Borden. 1996. Distribution and impact of *Leptoglossus occidentalis* Heidemann (Hemiptera: Coreidae) in seed orchards in British Columbia. *The Canadian Entomologist* 128:1065-1076.

- Brooks, A.R. and L.A. Kelton. 1967. Aquatic and semiaquatic Heteroptera of Alberta, Saskatchewan, and Manitoba (Hemiptera). *Memoirs of the Entomological Society of Canada* 51:1-92.
- De Clercq, P. 2000. Predaceous stinkbugs (Pentatomidae: Asopinae). Pp. 737-789. *In* Schaefer, C.W. and A.R. Panizzi (eds.). *Heteroptera of Economic Importance*. CRC Press, Boca Raton, FL.
- Drake, C.J. and H.M. Harris 1934. The Gerrinae of the Western Hemisphere (Hemiptera). *Annals of the Carnegie Museum* 23:179-240.
- Drake, C.J. and J.D. Lattin. 1963. American species of the lacebug genus *Acalypta* (Hemiptera: Tingidae). *Proceedings of the United States National Museum* 115:331-345.
- Foldessy, M. 2006. The presence of *Leptoglossus occidentalis* Heidemann, 1910 (Heteroptera: Coreidae) in North-east Hungary. *Folia Historico Naturalia Musei Matraensis* 30:203-204.
- Froeschner, R.C. 1960. Cydnidae of the Western Hemisphere. *Proceedings of the United States National Museum* 111:337-680.
- Gall, W. 1992. Further eastern range extensions of *Leptoglossus occidentalis*. *Great Lakes Entomologist* 25:159-171.
- Gogala, A. 2003. Listonozka (*Leptoglossus occidentalis*) ze v Sloveniji (Heteroptera: Coreidae). *Acta Entomologica Slovenica* 11:189-190.
- Harris, H.M. 1928. A monographic study of the hemipterous family Nabidae as it occurs in North America. *Entomologica Americana* 9:1-97.
- Harris, H.M. and W.E. Shull. 1944. A preliminary list of Hemiptera of Idaho. *Iowa State College Journal of Science* 18:199-208.
- Hedlin, A.F., H.O. Yates, D.C. Tovar, B.H. Ebel, T.W. Kroeber and E.P. Merkel. 1981. Cone and Seed Insects of North American Conifers. Canadian Forestry Services, USDA Forest Service and Secretaria de Agricultura y Recursos Hidraulicos, Mexico. 122 pp.
- Henry, T.J. 1979. Review of the *Ceratopsus lutescens* group, with descriptions of seven new species from the eastern United States (Hemiptera: Miridae). *Proceedings of the Entomological Society of Washington* 81:401-423.
- Henry, T.J. and J.D. Lattin 1988. Family Nabidae Costa, 1853. The Damsel Bugs. Pp. 508-520. *In* T.J. Henry and R.C. Froeschner (eds.). *Catalog of the Heteroptera, or True Bugs, of Canada and the Continental United States*. E.J. Brill, Leiden.
- Henry, T.J. and A.G. Wheeler, Jr. 1988. Family Miridae Hahn, 1833 (= Capsidae Burmeister, 1835). The Plant Bugs. Pp. 251-507. *In* T.J. Henry and R.C. Froeschner (eds.). *Catalog of the Heteroptera, or True Bugs, of Canada and the Continental United States*. E.J. Brill, Leiden.
- Henry, T.J., A.G. Wheeler, Jr. and W.E. Steiner, Jr. 2008. First North American records of *Amphiareus obscuriceps* (Poppius) (Hemiptera: Heteroptera: Anthocoridae), with a discussion of dead-leaf microhabitats. *Proceedings of the Entomological Society of Washington* 110:402-416.
- Hilpold, A. 2005. Neu fuer Suedtirol: *Leptoglossus occidentalis* Heidemann, 1910 (Heteroptera, Coreidae). *Gredleriana* 5:358.
- Hoebcke, E.R. and A.G. Wheeler, Jr., 1982. *Rhopalus (Brachycarenum) tigrinus*, recently established in North America, with a key to the genera and species of Rhopalidae in eastern North America (Hemiptera: Heteroptera). *Proceedings of the Entomological Society of Washington* 84:213-224.
- Hungerford, H.B. 1933. The genus *Notonecta* of the world. *University of Kansas Science Bulletin* 21:5-195.
- Hungerford, H.B. 1948. The Corixidae of the Western Hemisphere (Hemiptera). *University of Kansas Science Bulletin* 32:5-827.
- Jansson, A. 2002. New records of Corixidae (Heteroptera) from northeastern USA and eastern Canada, with one new synonymy. *Entomologica Fennica* 13:85-88.
- Kelton, L.A. 1978. The Anthocoridae of Canada and Alaska. Heteroptera: Anthocoridae. *The Insects and Arachnids of Canada. Part 4. Agriculture Canada Research Branch Publication* 1639: 101 pp.
- Kelton, L.A. 1980. The plant bugs of the Prairie Provinces of Canada. *The Insects and Arachnids of Canada. Part 8. Agriculture Canada Research Branch Publication* 1703: 408 pp.
- Kenner, R.D. and K.M. Needham. 2004. New waterboatmen records for Western Canada (Hemiptera: Corixidae). *Journal of the Entomological Society of British Columbia* 101:147-148.
- Kerzhner, I.M. 1963. Beitrag zur Kenntnis der Unterfamilie Nabinae (Heteroptera: Nabidae). *Acta Entomologica Musei Nationalis Prague* 35:5-61.
- Kerzhner, I.M. 1993. New and little-known Nabidae from North America (Heteroptera). *Zoosystematica Rossica* 1 (1992):37-45.
- Knight, H.H. 1920. New and little-known species of *Phytocoris* from the eastern United States (Heteroptera-Miridae). *Bulletin of the Brooklyn Entomological Society* 15:49-66.
- Knight, H.H. 1921. Monograph of the North American species of *Deraeocoris* (Heteroptera: Miridae). *University of Minnesota Agricultural Experimental Station Technical Bulletin* 1:75-210.
- Knight, H.H. 1927. Descriptions of seven new species of the genus *Orthotylus* Fieber (Hemiptera, Miridae).

- The Canadian Entomologist 59:176-181.
- Knight, H.H. 1941. The plant bugs, or Miridae, of Illinois. Illinois Natural History Survey Bulletin 22:1-234.
- Knight, H.H. 1968. Taxonomic review: Miridae of the Nevada test site and the western United States. Brigham Young University Science Bulletin (Biology Series) 9(3):1-282.
- Knight, H.H. 1974. A key to species of *Phytocoris* Fallen belonging to the *Phytocoris junceus* Knegt. group of species (Hemiptera, Miridae). Iowa State Journal of Research 49:123-135.
- Koeber, T.W. 1963. *Leptoglossus occidentalis* (Hemiptera, Coreidae), a newly discovered pest of coniferous seed. Annals of the Entomological Society of America. 56:229-234.
- Krugman, S.L. 1969. Effect of cone feeding by *Leptoglossus occidentalis* on ponderosa pine seed development. Forest Science 15:104-111.
- Larochelle, A. 1984. Les Punaises Terrestres (Hemipteres: Géocorises) du Quebec. Fabriques Supplement 3: 513 pp.
- Lattin, J.D. 1999. Bionomics of the Anthocoridae. Annual Review of Entomology 44:207-231.
- Lattin, J.D. 2000. Minute pirate bugs (Anthocoridae). Pp. 607-637. In C.W. Schaefer and A.R. Panizzi (eds.). Heteroptera of Economic Importance. CRC Press, Boca Raton, FL.
- Lattin, J.D. 2004. *Solenonotus angustatus* Poppius, a synonym of *Dufouriellus ater* Fieber (Hemiptera: Heteroptera: Anthocoridae). Proceedings of the Entomological Society of Washington 106:722-724.
- Lattin, J.D. 2005. *Dufouriellus ater* (Puton), *Macrotrachelia nigronitens* (Stål) and *Xylocoris (Arrostelus) flavipes* (Reuter) (Hemiptera: Heteroptera: Cimicoidea: Anthocoridae): first records from the Hawaiian Islands. Proceedings of Entomological Society of Washington 107:466-468.
- Lattin, J.D. 2007a. The Lasiophilidae, Lyctocoridae, and Anthocoridae (Hemiptera: Heteroptera) of the Hawaiian Islands: native or introduced? Proceedings of Entomological Society of Washington 109:75-80.
- Lattin, J.D. 2007b. The non-indigenous Lyctocoridae and Anthocoridae (Hemiptera: Heteroptera; Cimicoidea) of America north of Mexico. Proceedings of Entomological Society of Washington 109:366-376.
- Lewis, T.M., D.R. Horton and D.A. Broers. 2005. New state and United States records for Anthocoridae (Hemiptera: Heteroptera). Pan-Pacific Entomologist 81:59-67.
- Lindskog, P. 1981. On the identity of *Saldula laticollis* (Reuter) (Heteroptera: Saldidae). Pan-Pacific Entomologist 57:321-326.
- Marshall, S.A. 1991. A new Ontario record of a seed eating bug (Hemiptera: Coreidae) and other examples of the role of regional insect collections in tracking changes in Ontario's fauna. Proceedings of the Entomological Society of Ontario 122:109-111.
- Matsuda, R. 1977. The Aradidae of Canada. Hemiptera: Aradidae. The Insects and Arachnids of Canada. Part 3. Research Branch Canada Department of Agriculture Publication. 1634: 116 pp.
- Maw, H.E.L, R.G. Foottit, K.G.A. Hamilton and G.G.E. Scudder. 2000. Checklist of the Hemiptera of Canada and Alaska. NRC Research Press, Ottawa. 220 pp.
- McDonald, F.J.D. 1986. Revision of *Cosmopepla* Stål (Hemiptera: Pentatomidae). Journal of the New York Entomological Society 94:1-15.
- McPherson, J.E. 1982. The Pentatomoidea (Hemiptera) of northeastern North America, with emphasis on the fauna of Illinois. Southern Illinois University Press, Carbondale and Edwardsville. 240 pp.
- McPherson, J.E., R.J. Packauskas, S.F. Taylor and M.F. O'Brien. 1990. Eastern range extension of *Leptoglossus occidentalis*, with a key to *Leptoglossus* species of America north of Mexico (Heteroptera: Coreidae). Great Lakes Entomologist 23:99-104.
- Mitchell, P.L. 2000. Leaf-footed bugs (Coreidae). Pp 337-403. In C.W. Schaefer and A.R. Panizzi (eds.). Heteroptera of Economic Importance. CRC Press, Boca Raton, FL.
- Moulet, P. 2006. Un nouveau Coreide en France: *Leptoglossus occidentalis* Heidemann, 1910 (Heteroptera Coreidae). Entomologiste, Paris 62:183-184.
- Paiero, S.M., S.A. Marshall and K.G.A. Hamilton. 2003. New records of Hemiptera from Canada and Ontario. Journal of the Entomological Society of Ontario 134:115-129.
- Panizzi, A.R., J.E. McPherson, D.G. James, M. Javahery and McPherson, R.M. 2000. Stink Bugs (Pentatomidae). Pp. 421-474. In Schaefer, C.W. and A.R. Panizzi (eds.). Heteroptera of Economic Importance. CRC Press, Boca Raton, FL.
- Polhemus, D.A. 1994. An annotated checklist of the plant bugs of Colorado (Heteroptera: Miridae). Pan-Pacific Entomologist 70:122-147.
- Polhemus, J.T. 1985. Shore Bugs (Heteroptera, Hemiptera; Saldidae). A world overview and taxonomy of Middle American forms. The Different Drummer, Englewood, Co. 252 pp.
- Polhemus, J.T. 1988. Family Saldidae Amyot and Serville, 1843. The Shore Bugs. Pp. 665-681. In T.J. Henry and R.C. Froeschner (eds.). Catalog of the Heteroptera, or True Bugs, of Canada and the Continental United States. E.J. Brill, Leiden.
- Polhemus, J.T. and H.C. Chapman. 1979. Family Saldidae / Shore Bugs. Pp. 16-33. In A.S. Menke (ed.).

- The Semiaquatic and Aquatic Hemiptera of California (Heteroptera: Hemiptera). Bulletin of the California Insect Survey. Volume 21.
- Rabitsch, W. and E. Heiss. 2005. *Leptoglossus occidentalis* Heidemann, 1910, eine amerikanische Adventivart auch in Oesterreich aufgefunden (Heteroptera: Coreidae). Berichte des Naturwissenschaftlich Medizinischen Vereins in Innsbruck 92:131-135.
- Ribes, J. and E.-B. Oleguer. 2005. *Leptoglossus occidentalis* Heidemann, 1910, hemipter nearctic trobat a Catalunya (Hemiptera: Heteroptera: Coreidae). Sessio Conjunta d'Entomologia ICHN SCL 13 (2003):47-50.
- Ridge-O'Connor, G.E. 2001. Distribution of the western conifer seed bug, *Leptoglossus occidentalis* Heidemann (Heteroptera: Coreidae) in Connecticut and parasitism by a tachinid fly, *Trichopoda pennipes* (F.) (Diptera: Tachinidae). Proceedings of the Entomological Society of Washington 103:364-366.
- Roch, J.-F. 2007. Liste des punaises du Québec et des Régions adjacentes (Hemiptera: Heteroptera). Entomofauna du Québec Document Faunique 27: 42 pp.
- Rolston, L.H. 1973. The genus *Meneclis* Stål (Hemiptera: Pentatomidae). Journal of the New York Entomological Society 80:234-237.
- Rolston, L.H. 1983. A revision of the genus *Acrosternum* Fieber, subgenus *Chinavia* Orian, in the Western Hemisphere (Hemiptera: Pentatomidae). Journal of the New York Entomological Society 91:97-176.
- Ruth, D.S., G.E. Miller and J.R. Sutherland. 1982. A guide to common insect pests and diseases in spruce seed orchards in British Columbia. Canadian Forestry Service, Environment Canada. BC-X-231.
- Schaefer, C.W. and N.P. Chopra. 1982. Cladistic analysis of the Rhopalidae, with a list of food plants. Annals of the Entomological Society of America 75:224-233.
- Schaefer, C.W. and P.L. Mitchell. 1983. Food plants of the Coreidae (Hemiptera: Heteroptera). Annals of the Entomological Society of America 76:591-615.
- Schaffner, J.C. 1967. The occurrence of *Theognis occidentalis* in the midwestern United States (Heteroptera: Coreidae). Journal of the Kansas Entomological Society 40:141-142.
- Schowalter, T.D. and J.M. Sexton. 1990. Effect of *Leptoglossus occidentalis* (Heteroptera: Coreidae) on seed development of Douglas fir at different times during the growing season in western Oregon. Journal of Economic Entomology 83:1485-1486.
- Schuh, R.T. 2000a. Revision of *Oligotylus* Van Duzee with descriptions of ten new species from western North America and comments on *Lepidargyrus* in the Nearctic (Heteroptera: Miridae: Phylinae: Phylini). American Museum Novitates 3300: 44 pp.
- Schuh, R.T. 2000b. Revision of the North American plant bug genus *Megalopsallus* Knight, with description of eight new species from the west (Heteroptera: Miridae: Phylinae). American Museum Novitates 3305: 69 pp.
- Schuh, R.T. 2001. Revision of New World *Plagiognathus* Fieber, with comments on the Palearctic fauna and the description of a new genus (Heteroptera: Miridae: Phylinae). Bulletin of the American Museum of Natural History 266: 267 pp.
- Schuh, R.T. 2004a. Revision of *Tuxedo* Schuh (Hemiptera: Miridae: Phylinae). American Museum Novitates 3435: 26 pp.
- Schuh, R.T. 2004b. Revision of *Europiella* Reuter in North America, with the description of a new genus (Heteroptera: Miridae: Phylinae). American Museum Novitates 3463: 58 pp.
- Schuh, R.T. and M.D. Schwartz. 1988. Revision of the New World Pilophorini (Heteroptera: Miridae: Phylinae). Bulletin of the American Museum of Natural History 187:101-201.
- Schuh, R.T. and M.D. Schwartz. 2004. New genera, new species, new synonyms, and new combinations in North American and Caribbean Phylinae (Heteroptera: Miridae). American Museum Novitates 3436: 36 pp.
- Schuh, R.T. and M.D. Schwartz. 2005. Review of North American *Chlamydatus* Curtis species, with new synonymy and the description of two new species (Heteroptera: Miridae: Phylinae). American Museum Novitates 3471: 55 pp.
- Schuh, T. 1967. The shore bugs (Hemiptera: Saldidae) of the Great Lakes region. Contributions of the American Entomological Institute 2(2): 35 pp.
- Schwartz, M.D. and G.G.E. Scudder 2001. Miridae (Heteroptera) new to Canada, with some taxonomic changes. Journal of the New York Entomological Society 108 (2000):248-267.
- Schwartz, M.D. and G.G.E. Scudder. 2003. Seven new species of Miridae (Heteroptera) from British Columbia and Alaska and synonymy of *Adelphocoris superbus* (Uhler). Journal of the New York Entomological Society 111:65-95.
- Schwartz, M.D. and G.M. Stonedahl. 2004. Revision of *Phoenicocoris* Reuter with descriptions of three new species from North America and a new genus from Japan (Heteroptera: Miridae: Phylinae). American Museum Novitates 3464: 55 pp.
- Scudder, G.G.E. 1962. The Ischnorhynchinae of the world (Hemiptera: Lygaeidae). Transactions of the

- Royal Entomological Society of London 114:163-194.
- Scudder, G.G.E. 1977. An annotated checklist of the aquatic and semiaquatic Hemiptera (Insecta) of British Columbia. *Syesis* 10:31-38.
- Scudder, G.G.E. 1997. True Bugs (Heteroptera) of the Yukon. Pp. 241-336. In H.V. Danks and J.A. Downes (eds.). *Insects of the Yukon. Biological Survey of Canada (Terrestrial Arthropods)*, Ottawa.
- Scudder, G.G.E. 2000. Heteroptera (Hemiptera: Prosorrhyncha) new to Canada. Part. 1. *Journal of the Entomological Society of British Columbia* 97:51-56.
- Scudder, G.G.E. 2004. Heteroptera (Hemiptera: Prosorrhyncha) new to Canada. Part 2. *Journal of the Entomological Society of British Columbia* 101:125-129.
- Scudder, G.G.E. 2007. Two alien Heteroptera (Hemiptera) new to Canada. *Journal of the Entomological Society of British Columbia* 104:81-84.
- Scudder, G.G.E. and S.S. Duffey. 1972. Cardiac glycosides in the Lygaeinae (Hemiptera: Lygaeidae). *Canadian Journal of Zoology* 50:35-42.
- Scudder, G.G.E. and R.G. Footitt 2006. Alien true bugs (Hemiptera: Heteroptera) in Canada: composition and adaptations. *The Canadian Entomologist* 138:24-51.
- Scudder, G.G.E. and M.D. Schwartz 2001. The genus *Leptopterna* Fieber (Heteroptera: Miridae: Stenodemini) in North America. *Proceedings of the Entomological Society of Washington* 103(4):797-806.
- Slater, A. 1988. A new genus and two new species of Lygaeinae (Heteroptera: Lygaeidae). *Journal of the Kansas Entomological Society* 61:308-316.
- Slater, J.A. 1954. Notes on the genus *Labops* Burmeister in North America, with the description of three new species (Hemiptera: Miridae). *Bulletin of the Brooklyn Entomological Society* 49:57-65, 89-94.
- Slater, J.A. and R.M. Baranowski 1978. How to know the true bugs (Hemiptera-Heteroptera). Wm. C. Brown Company Publishers, Dubuque, Iowa. 256 pp.
- Stock, M.W. and J.D. Lattin. 1976. Biology of intertidal *Saldula palustris* (Douglas) on the Oregon coast (Heteroptera: Saldidae). *Journal of the Kansas Entomological Society* 49:313-326.
- Strong, W.B. 2006. Seasonal changes in seed reduction in Lodgepole pine cones caused by feeding of *Leptoglossus occidentalis* (Hemiptera: Coreidae). *The Canadian Entomologist* 138:888-896.
- Strong, W.B., S.L. Bates and M.U. Stoehr. 2001. Feeding by *Leptoglossus occidentalis* Heidemann (Hemiptera: Coreidae) reduces seed set in lodgepole pine. *The Canadian Entomologist* 133:857-865.
- Sweet, M.H. 1964. The biology and ecology of the Rhyparochrominae of New England (Heteroptera: Lygaeidae). Part II. *Entomologica Americana (new series)* 44:1-201.
- Taylor, S.J., G. Tescari and M. Villa. 2001. A nearctic pest of Pinaceae accidentally introduced into Europe: *Leptoglossus occidentalis* (Heteroptera: Coreidae) in northern Italy. *Entomological News* 112:101-103.
- Tescari, G. 2004. First record of *Leptoglossus occidentalis* (Heteroptera: Coreidae) in Croatia. *Entomologia Croatica* 8:73-75.
- Thomas, D.B. 1992. Taxonomic synopsis of the Asopine Pentatomidae (Heteroptera) of the Western Hemisphere. *The Thomas Say Foundation Volume* 16: 156 pp.
- Tinerella, P.P. and R.W. Gundersen 2005. The water boatmen (Insecta: Heteroptera: Corixidae) of Minnesota. *Schafer-Post Series: North Dakota Insects Publication* 23: 119 pp.
- Van Duzee, E.P. 1916. Checklist of the Heteroptera (excepting the Aphididae, Aleurodidae and Coccidae) of America, north of Mexico. *New York Entomological Society, New York*. 111 pp.
- Van Duzee, E.P. 1917. Catalogue of the Hemiptera of America north of Mexico, excepting the Aphididae, Coccidae and Aleurodidae. *University of California Publications, Technical Bulletins, Entomology* 2: 902 pp.
- Vinokurov, N.N. 1988. *Heteroptera of Yakutia*. Oxonian Press Pvt. Ltd., New Delhi, Calcutta. 328 pp.
- Wheeler, Jr., A.G., and T.J. Henry 1992. A Synthesis of the Holarctic Miridae (Heteroptera): Distribution, Biology, and Origin, with Emphasis on North America. *The Thomas Say Foundation Volume* 15: 282 pp.
- Wheeler, Jr., A.G., T.J. Henry and E.R. Hoebeke 2006. Palearctic plant bugs (Hemiptera, Miridae) in Newfoundland, Canada: First North American records for *Phytocoris longipennis* Flor and *Pilophorus cinnamopterus* (Kirschbaum), new records of eight other species, and review of previous reported species. *Denisia* 19 (2006):997-1014.
- Wheeler, Jr., A.G., T.J. Henry and T.L. Mason, Jr. 1983. An annotated list of the Miridae of West Virginia (Hemiptera-Heteroptera). *Transactions of the American Entomological Society* 109:127-159.
- Wheeler, Jr., A.G. and E.R. Hoebeke. 2004. New records of Palearctic Hemiptera (Sternorrhyncha, Cicadomorpha, Heteroptera) in the Canadian maritime provinces. *Proceedings of the Entomological Society of Washington*. 106:298-304.
- Wright, B. 1989. The fauna of Sable Island. *Nova Scotia Museum Curatorial Report* 68: 93 pp.